

ORIGINAL FILE COPY

2

AD-A230 758

OFFICE OF NAVAL RESEARCH

Grant N00014-90-J-1193

TECHNICAL REPORT No. 39

Cluster Calculation of  $\text{CuO}_2$  in High- $T_c$  Superconductors

by

H. R. Lee, Ki-Soo Sohn and Thomas F. George

Prepared for publication

in

*The Challenges to Advanced New Materials*

*12th Kyungpook National University International Seminar Proceedings*

Taegu, Korea, 1990

Departments of Chemistry and Physics  
State University of New York at Buffalo  
Buffalo, New York 14260

January 1991

Reproduction in whole or in part is permitted for any purpose of the  
United States Government.

This document has been approved for public release and sale;  
its distribution is unlimited.

DTIC  
S ELECTE D  
JAN 11 1991  
E

91 1 10 022

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

Fc Approved  
OML No. 0704-0188

a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
1a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
1b. DECLASSIFICATION / DOWNGRADING SCHEDULE			
1. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/91/TR-39		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
5a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION	
5c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260		7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Grant N00014-90-J-1193	
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO.	PROJECT NO.
		TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Cluster Calculation of CuO <sub>2</sub> in High-T <sub>c</sub> Superconductors			
12. PERSONAL AUTHOR(S) H. R. Lee, Ki-Soo Sohn and Thomas F. George			
13a. TYPE OF REPORT	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) January 1991	15. PAGE COUNT 12
16. SUPPLEMENTARY NOTATION Prepared for publication in <i>The Challenges to Advanced New Materials</i> , 15th Kyungpook National University International Seminar Proceedings Taegu, Korea, 1990			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		HIGH-TEMPERATURE SUPERCONDUCTIVITY	
		CuO <sub>2</sub> PLANE	
		EXTENDED HUBBARD HAMILTONIAN	
		CLUSTER CALCULATION	
		MAGNETIC SUSCEPTIBILITY	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
<p>The eigenvalues of a small cluster in the CuO<sub>2</sub> plane are obtained by exactly diagonalizing the extended Hubbard Hamiltonian. This eigenenergy spectrum is used for the grand partition function. The magnetic and thermodynamic quantities such as the magnetic susceptibility and specific heat are calculated with different occupation numbers and discussed in terms of the excited states of the system.</p>			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson		22b. TELEPHONE (Include Area Code) (202) 696-4410	22c. OFFICE SYMBOL

*The Challenges to Advanced New Materials*  
*Twelfth Kyungpook National University International Seminar Proceedings*  
*Taegu, Korea, 1990*

Cluster Calculation of  $\text{CuO}_2$  in High- $T_c$  Superconductors

H. R. Lee and Ki-Soo Sohn

Department of Physics  
Kyungpook National University  
Taegu, 702-701 KOREA

Thomas F. George

*Departments of Physics & Astronomy and Chemistry*  
*239 Fronczak Hall*  
*State University of New York at Buffalo*  
*Buffalo, New York 14260*

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

**Abstract**

The eigenvalues of a small cluster in the  $\text{CuO}_2$  plane are obtained by exactly diagonalizing the extended Hubbard Hamiltonian. This eigenenergy spectrum is used for the grand partition function. The magnetic and thermodynamic quantities such as the magnetic susceptibility and specific heat are calculated with different occupation numbers and discussed in terms of the excited states of the system.

PACS Nos: 25.12.75.30.-m, 74.30.Ek

**I. Introduction**

Since the discovery of superconductivity in oxide compounds [1], many theoretical and experimental studies were carried out for the pairing mechanism on these compounds. For most crystalline solids, the electronic properties are well described by the band theory in



terms of a one-body Hamiltonian. But it is now generally believed that the superconducting oxide compound do not fit this description. In this case, the excitations of these systems are totally beyond the description of the band-structure approximation. Therefore, in this paper, we study the magnetic and thermodynamic properties in a small cluster with an appropriate extended Hubbard Hamiltonian, because those properties mainly depend on the excitation spectrum of the system.

Even though the high-temperature oxide superconductors have some similarities with conventional superconductors, there seems to be a fundamental difference for pairing mechanism. In conventional superconductors, the attraction between two electrons arises from the phonon exchange, where the retarded phonon attraction is able to overcome the screened Coulomb repulsion between two electrons. However it is very hard to consider a similar mechanism for high-temperature oxide superconductors with short coherent length  $\xi_c \sim d$ , where  $d$  is the interparticle distance between electrons. The wide range of energy gap instead of a sharp gap observed in most tunneling measurements [2] seems due to the fluctuation effect, which is expected to be considerably large due to the short coherent length. More recently discovered n-type superconductors[3], which have similar crystal structures to the p-type, provide additional information about these compounds: (1) the superconducting pairs occur in purely two dimensions, and (2) there must be electron-hole symmetry for whatever band is used in the Hamiltonian.

## II. Theory

The central physics in the theory of high-temperature superconductivity centers on which effective Hamiltonian one should take to describe the electrons in the  $\text{CuO}_2$  plane. The high- $T_c$  compounds can be explained well by the extended Hubbard Hamiltonian. In previous papers[4], we have pointed out two important physical features of the pairing of electrons and holes in n-type and p-type oxide superconductors, respectively: (i) the vacuum level must be defined as  $(d^9p^6)$  for the copper and oxygen sites instead of the closed orbit configuration  $(d^{10}p^6)$  used by other [5]. x-ray absorption spectra show that the charge carriers (electrons) are mostly located on the copper sites ( $\text{Cu}^{2+}$ ) in n-type superconductors[6], while the charge

carriers (holes) are located on oxygen sites ( $O^{1-}$ ) in p-type superconductors [7]. By this observation, it is appropriate to define the vacuum level to be ( $d^9p^6$ ), which is an undoped configuration for both types and plays a critical role for fixing the electron and the hole energy levels. (2) Since holes are not true particles but are the empty states which represent the absence of electrons, we have to be more careful in setting up the Hubbard Hamiltonian. Generally the on-site Coulomb repulsion is introduced in the Hubbard Hamiltonian when two electrons occupy the same orbit. This cannot be the case for the hole states, if one electron is removed from two electron states of the given orbit, i. e., one hole is created, the on-site Coulomb repulsion disappears.

In other words, the on-site Coulomb repulsion, which is usually the largest parameter in the Hubbard Hamiltonian, appears whenever a hole is created from the given closed shell orbit instead of when two holes are created.

Considering the above two features, the extended Hubbard Hamiltonian can be written as

$$\begin{aligned}
 [4] \quad H = & t_{pd} \sum_{\langle i,j \rangle} (d_{jo}^h p_{jo} + d_{jo}^e p_{jo}^* + \text{h.c.}) \\
 & + \sum_{i,o} (E_d + U_d) d_{io}^e d_{io}^e + \sum_{i,o} E_d d_{io}^h d_{io}^h \\
 & - \sum_{i,o} (E_p + U_p) p_{io}^e p_{io}^e - \sum_{i,o} U_p n_{io} n_{io} \\
 & - \sum_{i,j} U_{pd} n_{io}^e n_{jo}^h,
 \end{aligned}
 \quad (1)$$

where  $t_{pd}$  is the hopping integral between neighboring Cu and O sites, and the  $d$ 's and  $p$ 's are second quantization operators related to the Cu and O sites, respectively. The superscripts h and e on the Cu site operators represent the holes  $Cu^{1+}(d^9)$  and electrons  $Cu^{2+}(d^9)$ , explicitly. The prime on a summation indicates  $i \neq j$ , and  $\langle i, j \rangle$  indicates nearest neighbors between the Cu and O sites.  $E_d$  and  $E_p$  are the diagonal kinetic energies of the 3d and 2p orbitals, respectively. The electron-hole pair creation potential  $U_{pd}$  in the last term of the Hamiltonian is introduced to account for the largest valence charge difference when an electron on an O site moves to a neighboring Cu site, i. e.,  $Cu^{2+}(d^9) + O^{1-}(p^6) \rightarrow Cu^{1+}(d^9) + O^{2-}(p^5)$ . As mentioned earlier, the on-site Coulomb repulsions  $U_d$  and  $U_p$  appear when an electron is created on a Cu site or a hole is created on an O site. With the condition  $E_d + U_d > E_p + U_p$ , the

undoped  $\text{La}_2\text{CuO}_4$  and  $\text{Nb}_2\text{CuO}_4$  nominally consist of  $\text{Cu}^{2+}$  and  $\text{O}^{2-}$ , and the holes introduced by doping with the divalent Sr in  $\text{La}_2\text{CuO}_4$  occupy the oxygen orbitals in p-type superconductors, while the electrons introduced by doping with the tetravalent Th in  $\text{Nd}_2\text{CuO}_4$  occupy the copper orbitals in n-type superconductors. Clearly this observation is due to the presence or absence of electrons in the antibonding states of  $\text{Cu}(d_{x^2-y^2})$  and  $\text{O}(p_x \text{ or } p_y)$  orbitals.

We believe this hybridization is strongly related to the quick disappearance of the antiferromagnetic background on the Cu sites with a small amount of doping. Even though the direct hopping integral ( $t_{\infty}$ ) between neighboring oxygen sites is not negligible, the direct oxygen band is supposed to be much lower than the antibonding states of the Cu and O orbitals[8] due to the condition  $E_d + U_d > E_o + U_o$ . Therefore, the direct oxygen hopping integral is not included in the Hamiltonian.




Fig 1. The cluster considered in the calculation includes two unit cells in the  $\text{CuO}_2$  square lattice. The copper sites and oxygen sites are represented by the circles and plus signs, respectively.

Based on the above arguments, we have proposed a local pairing mechanism and estimated the binding energy of two carriers in a previous paper[4]. In the present study, we examine the magnetic and thermodynamic quantities such as the magnetic susceptibility and specific heat by exact diagonalization on small two-dimensional clusters of two unit cells on a  $\text{CuO}_2$  lattice as shown in Fig. 1. By introducing the chemical potential  $\mu$  into the Hamiltonian

an, we can allow the number of electrons to vary in this system. In this case, all the second quantization operators in Eq.(1) can be changed to the true electron representation without the hole picture. Information on the interaction between the cluster and the rest of the system is also included in the chemical potential. The Hubbard Hamiltonian, Eq.(1), of this system is now modified in the presence of a uniform magnetic field as

$$\begin{aligned}
 H = & t_{pd} \sum_{\langle i,j \rangle} (d_{i0}^\dagger p_{j0} + \text{h.c.}) \\
 & + \sum_{i,b} E_d d_{i0}^\dagger d_{i,b} + \sum U_d n_{i0} n_{i,b} \\
 & + \sum_{i,b} E_p p_{i0}^\dagger p_{i,b} + \sum U_p n_{i0} n_{i,b} \\
 & + \sum_{\substack{\langle i,j \rangle \\ b,b'}} U_{pd} n_{i,b} n_{j,b'} - \mu \sum_{i,b} n_{i,b} - \gamma B \sum_i (n_{i0} n_{i,b'})
 \end{aligned} \quad (2)$$

where  $\gamma$  is the magnetic moment of the electron, and the uniform magnetic field  $B$  is perpendicular to the  $\text{CuO}_2$  plane.

The eigenvalues of the above Hamiltonian can be obtained by diagonalizing exactly in a small cluster system. The grand partition function  $Z$  can be written immediately with the eigenvalues. To calculate the magnetic and thermodynamic quantities, we have to first evaluate the chemical potential of the system for the given occupation numbers. The chemical potential can be determined by inverting the expression

$$\langle N \rangle = K_B T \left[ \frac{\partial \ln Z}{\partial \mu} \right]_{T, V} \quad (3)$$

in terms of the occupation number  $N$  and the other parameters in the Hamiltonian. The magnetic susceptibility  $\chi$  and the specific heat  $C_v$  can be calculated with a knowledge of the grand partition function and the chemical potential in the usual manner.

### III. Numerical Calculations and Discussion

Considering four possible states per site, the given clusters has  $4^N$  eigenvalues and eigenvectors. Since the Hamiltonian is isotropic in spin space, the total spin  $S$  and its  $z$ -com-

ponent  $S_z$  are good quantum numbers. Then the  $4^N \times 4^N$  Hamiltonian matrix can be reduced to many blocks with different number of occupations. In the following calculations, we have used these reported parameters, which are in the most acceptable range [8]:  $\Delta E = 1.5$  eV,  $t_{pd} = 1.2$  eV,  $U_d = 9.0$  eV,  $U_p = 6.0$  eV and two different values of (1)  $U_{pd} = 0.0$  eV, (2)  $U_{pd} = 1.5$  eV. In all cases, it is found that the ground state in the state is the state with minimal  $|S_z|$ . This result is due to the relatively large hopping integral  $t_{pd}$  and more degree of freedom with less  $|S_z|$ . The behavior of the magnetic and thermodynamic quantities in the cluster model can be understood from the excitation spectra. In Figs. 2 and 3, we have shown the magnetic susceptibility and specific heat curves of the two unit-cell model for  $N = 10$ , which is a vacuum configuration in our definition. The low-temperature peaks in these figures are due to the transition from the ground state of singlet spin configuration on two copper sites to the triplet spin configuration which is the first excited state. By contrast, these peaks disappear for  $N = 9$  and  $N = 8$ , which are shown in Figs. 4 through 7.




Fig 2. Magnetic susceptibility curves ( $x/2r^2$ ) for  $N = 10$  with (1)  $U_{pd} = 0.0$  eV and (2)  $U_{pd} = 1.5$  eV.






Fig. 3 Specific heat curve ( $C_v/N$ ) for  $N=10$  with (1)  $U_{pd}=0.0\text{eV}$  and (2)  $U_{pd}=1.5\text{eV}$ .




Fig. 4 Magnetic susceptibility curves ( $\chi/2r^2$ ) for  $N=9$  with (1)  $U_{pd}=0.0\text{ eV}$  and (2)  $U_{pd}=1.5\text{eV}$ .




Fig. 5 Specific heat curves ( $C_v/N$ ) for  $N=9$  with (1)  $U_d = 0.0\text{eV}$  and (2)  $U_d = 1.5\text{eV}$ .




Fig. 6 Magnetic susceptibility curves ( $x/2r^2$ ) for  $N=8$  with (1)  $U_d = 0.0\text{ eV}$  and (2)  $U_d = 1.5\text{eV}$ .




Fig. 7. Specific heat curves ( $C_v/N$ ) for  $N=8$  with (1)  $U_d = 0.0\text{eV}$  and (2)  $U_d = 1.5\text{eV}$ .

For more than half-filling cases ( $N>6$ ), the energy spectrum can roughly be divided into some groups with spacings of on-site Coulomb repulsion  $U_d$  or  $U_p$ , and the spacings between successive levels within the same group are expected to be on the order of  $t_{pd}^2/U_d$  or  $t_{pd}^2/U_p$ . As it turns out in our numerical calculations, the energy spacings between the ground state and the first excited state are greater than 0.5 eV for all cases except the vacuum configuration  $N=10$ . The energy difference between the ground state ( $S_z=0$ ) and first-excited state ( $S_z=1$ ) results in 30 meV~60 meV for  $N=10$  with various parameters in the Hamiltonian.

This energy difference gives the superexchange parameter  $J$  between neighboring copper sites in the Heisenberg representation,

$$H = J \sum_{\langle i, j \rangle} \vec{S}_i \cdot \vec{S}_j \quad (4)$$

which is a good approximation for an undoped material[9].  $J$  can be obtained alternatively by the fourth-order perturbation approximation to Eq.(1)[4],

$$J = \frac{4t_{pd}^4}{(\Delta E + U_d + U_{pd} + U_p)^2} \left[ \frac{1}{U_d} + \frac{2}{2\Delta E + 2U_d + 4U_{pd} - U_p} \right] \quad (5)$$

which has very good agreement with the above numerical result.

In Fig. 4, two magnetic susceptibility curves almost overlap and decreases monotonically as  $T$  increases because the ground level has a minimal  $S_z = 1/2$ , which is different from the case of the  $N = 8$  and  $N = 10$ . In Fig. 6, the magnetic susceptibility are increasing with increasing temperature as in Fig. 2, but no peaks appear under 1000 K due to the large excitation energy from the ground state, as explained earlier. These qualitatively different types of behavior of the magnetic susceptibility are merely caused by the smallness of the cluster. With the same explanation of large excitation energy from the ground state, the specific heat curves in Figs. 5 and 7 have no peaks in the low-temperature regime.

#### IV. Conclusions

The extended Hubbard Hamiltonian has been solved for clusters of two unit cells by an exact diagonalization. The ground states are always related to the minimal  $|S_z|$  configuration. The energy spectrum is grouped with the order of on-site coulomb repulsion  $U_s$  or  $U_d$ , and the successive levels in each group are separated at the order of  $t_{sd}^2/U_s$  or  $t_{sd}^2/U_d$ . But the first excited state in an undoped regime is the order of superexchange energy between nearest-neighbor copper sites, which agrees well with the result obtained by the fourth-order perturbation calculation. This small excitation energy contributes to the low-temperature peaks in the magnetic susceptibility and specific heat curves for  $N = 10$ . Due to the large excitation energy for the other configuration, no peaks appear in the magnetic susceptibility and specific heat curves in the low-temperature regime. From these results, we may conclude that the low-temperature physical anomalies appear only near the undoped regime. The magnetic susceptibility curve for the case of  $N = 9$  is essentially of the Curie form, monotonically decreasing as temperature increases. This is due to the odd number of electrons in a small cluster. We believe that this kind of behavior will be diminished if we increase the size of the cluster.

Finally, we would say that the results obtained in this calculation are all very stable with a wide range of parameters in the Hamiltonian.

Therefore, the qualitative nature of the results does not change within the acceptable parameter ranges, even though we have shown only two different values of  $U_{\infty}$ , which is supposed to be the most uncertain.

#### Acknowledgments

This research was supported by an award from the New York State Institute on Superconductivity in conjunction with the New York State Energy Research and Development Authority and by the Office of Naval Research.

#### References

1. J. G. Bednorz and K. A. Mueller, A. Phys. B64, 189(1986).
2. J. R. Kirtley, R. T. Collins, Z. Schlessinger, W. J. Gallagher, R. L. Sandstorm, T. R. Dinger and D. A. Chance, Phys. Rev. B35, 8846(1987) and Qiang Huang, J. F. Zasadzinski, K. E. Gray, J. Z. Liu and H. Claus, Phys. Rev. B40, 9366(1989), and the references therein.
3. Y. Tokura, H. Takagi and S. Uchida, Nature 337, 345(1989).
4. H. R. Iee, T. F. George and A. Langner, J. Phys.: Condens. Matter submitted; H. R. Iee and T. F. George, in Superconductivity and Applications, ed. by H. S. Kwok, Y. H. Kao and D. T. Shaw(Plenum, New York, 1989), in press.
5. E. B. Stechel and D. R. Jennison, Phys. Rev. B38, 4(1988) and V. J. Emery and G. Reifer, Phys. Rev. B38, (1988).
6. J. M. Tranquada, S. M. Heald, A. R. Moodenbaugh, G. Liang and J. Croft, Nature 37, 720 (1989).
7. J. M. Tranquada, S. M. Heald and A. R. Moodenbaugh, Phys. Rev. B36, 5263(1987).
8. J. M. Tranquada, S. M. Heald and A. R. Moodenbaugh, Phys. Rev. B36, 5263(1987).
9. A. K. McMahan, R. M. Martin and S. Satpathy, Phys. Rev. B38, 6650(1988); D. E. Ramaker, N. H. Turner, J. S. Murday, L. F. Toth, M. Osofsky and F. L. Hutson, Phys. Rev. B36, 5672(1987).
10. R. L. Greene, H. Maletta, T. S. Plaskett, J. G. Bednorz and K. A. Mueller, Solid State Commun. 63, 379(1987); K. Yamada, E. Kudo, Y. Endoh, Y. Hidaka, M. Oda, M. Suzuki and T. Murakami, ibid., 64, 753(1987).

FY90 Abstracts Distribution List for Solid State & Surface Chemistry

Professor John Baldeschwieler  
Department of Chemistry  
California Inst. of Technology  
Pasadena, CA 91125

Professor John Eyler  
Department of Chemistry  
University of Florida  
Gainesville, FL 32611

Dr. Sylvia Johnson  
SRI International  
333 Ravenswood Avenue  
Menlo Park, CA 94025

Professor Paul Barbara  
Department of Chemistry  
University of Minnesota  
Minneapolis, MN 55455-0431

Professor James Garvey  
Department of Chemistry  
State University of New York  
Buffalo, NY 14214

Dr. Zakya Kafafi  
Code 6551  
Naval Research Laboratory  
Washington, DC 20375-5000

Dr. Duncan Brown  
Advanced Technology Materials  
520-D Danury Rd.  
New Milford, CT 06776

Professor Steven George  
Department of Chemistry  
Stanford University  
Stanford, CA 94305

Professor Larry Kesmodel  
Department of Physics  
Indiana University  
Bloomington, IN 47403

Professor Stanley Bruckenstein  
Department of Chemistry  
State University of New York  
Buffalo, NY 14214

Professor Tom George  
Dept. of Chemistry and Physics  
State University of New York  
Buffalo, NY 14260

Professor Max Lagally  
Dept. Metal. & Min. Engineerin  
University of Wisconsin  
Madison, WI 53706

Professor Carolyn Cassady  
Department of Chemistry  
Miami University  
Oxford, OH 45056

Dr. Robert Hamers  
IBM T.J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, N Y 10598

Dr. Stephen Lieberman  
Code 522  
Naval Ocean Systems Center  
San Diego, CA 92152

Professor R.P.H. Chang  
Dept. Matls. Sci. & Engineering  
Northwestern University  
Evanston, IL 60208

Professor Charles Harris  
Department of Chemistry  
University of California  
Berkeley, CA 94720

Professor M.C. Lin  
Department of Chemistry  
Emory University  
Atlanta, GA 30322

Professor Frank DiSalvo  
Department of Chemistry  
Cornell University  
Ithaca, NY 14853

Professor John Hemminger  
Department of Chemistry  
University of California  
Irvine, CA 92717

Professor Fred McLafferty  
Department of Chemistry  
Cornell University  
Ithaca, NY 14853-1301

Dr. James Duncan  
Federal Systems Division  
Eastman Kodak Company  
Rochester, NY 14650-2156

Professor Leonard Interrante  
Department of Chemistry  
Rensselaer Polytechnic Institute  
Troy, NY 12181

Professor Horia Metiu  
Department of Chemistry  
University of California  
Santa Barbara, CA 93106

Professor Arthur Ellis  
Department of Chemistry  
University of Wisconsin  
Madison, WI 53706

Professor Roald Hoffmann  
Department of Chemistry  
Cornell University  
Ithaca, NY 14853

Professor Larry Miller  
Department of Chemistry  
University of Minnesota  
Minneapolis. MN 55455-0431

Professor Mustafa El-Sayed  
Department of Chemistry  
University of California  
Los Angeles, CA 90024

Professor Eugene Irene  
Department of Chemistry  
University of North Carolina  
Chapel Hill, NC 27514

Professor George Morrison  
Department of Chemistry  
Cornell University  
Ithaca, NY 14853

Professor Daniel Neumark  
Department of Chemistry  
University of California  
Berkeley, CA 94720

Professor Robert Whetten  
Department of Chemistry  
University of California  
Los Angeles, CA 90024

Professor David Ramaker  
Department of Chemistry  
George Washington University  
Washington, DC 20052

Professor R. Stanley Williams  
Department of Chemistry  
University of California  
Los Angeles, CA 90024

Dr. Gary Rubloff  
IBM T.J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, NY 10598

Professor Nicholas Winograd  
Department of Chemistry  
Pennsylvania State University  
University Park, PA 16902

Professor Richard Smalley  
Department of Chemistry  
Rice University  
P.O. Box 1892  
Houston, TX 77251

Professor Aaron Wold  
Department of Chemistry  
Brown University  
Providence, RI 02912

Professor Gerald Stringfellow  
Dept. of Matls. Sci.  
& Engineering  
University of Utah  
Salt Lake City, UT 84112

Professor Vicki Wysocki  
Department of Chemistry  
Virginia Commonwealth University  
Richmond, VA 23284-2006

Professor Galen Stucky  
Department of Chemistry  
University of California  
Santa Barbara, CA 93106

Professor John Yates  
Department of Chemistry  
University of Pittsburgh  
Pittsburg, PA 15260

Professor H. Tachikawa  
Department of Chemistry  
Jackson State University  
Jackson, MI 39217-0510

Professor William Unertl  
Lab. for Surface Sci.  
& Technology  
University of Maine  
Orono, ME 04469

Dr. Terrell Vanderah  
Code 3854  
Naval Weapons Center  
China Lake, CA 93555

Professor John Weaver  
Dept. of Chem. Eng. & Mat. Sci.  
University of Minnesota  
Minneapolis, MN 55455

Professor Brad Weiner  
Department of Chemistry  
University of Puerto Rico  
Rio Piedras, Puerto Rico 00931